

Slounase®

Hemocoagulase Injection



Product Name

Generic name: Hemocoagulase Injection

Trade name: Slounase®

Indication

To reduce bleeding or hemostasis in various medical conditions, such as surgery, internal medicine, obstetrics and gynecology, ophthalmology, otolaryngology, stomatology and other clinical departments of bleeding and bleeding diseases. It can also be used to prevent bleeding. Bleeding at the surgical site and after surgery can be avoided or reduced by applying hemocoagulase before surgery.

Dosage and administration

It can be administered by intravenous, intramuscular and subcutaneous injection. It can also be used as topical medication.

General bleeding: 1-2 units in adults; 0.3-0.5 units in children.

Emergency bleeding: 0.25-0.5 units intravenously immediately, and 1 unit intramuscularly at the same time.

Various surgical procedures: 1 unit intramuscularly the night before surgery, 1 unit intramuscularly 1 hour before surgery, 1 unit intravenously 15 minutes before surgery, and 1 unit intramuscularly daily for 3 days after surgery.

Hemoptysis: 1 unit subcutaneously every 12 hours, if necessary, add 1 unit intravenously at the beginning, preferably add 10ml of 0.9% NaCl solution for mixed injection.

Abnormal Bleeding: Double the dose, 1 unit intramuscularly at an interval of 6 hours until the bleeding has completely ceased.

Strength

1ml:1 Unit